

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

ronic

12

10

10.00

20

, 21 18-90 , (. 0.05)
15-30 d 讨论

30 , 5 ,

B

1 1 400IU/g

VAS

SS200

($\bar{x} \pm s$)

χ^2 . . 0.05

n n(%)

avar
eatn
e N

Hu

Chronic Wounds in a Specialized Wound Healing Department: A Retrospective Analysis of 107 Cases. *The International Journal of Lower Extremity Wounds*, 2015, 14(2):178-182

6 Bonomo RA, van Zile PS, Li Q, et al. Topical triple- antibiotic ointment as a novel therapeutic choice in wound management and infection prevention: a practical perspective. *Expert Review of Anti-infective Therapy*, 2007, 5(5):773-782

7 , , , . B , 2016, 37(9):2175-2176

8 , , , . B 2019-09-05

9 1(1):28-30

10 Ahmad FMF, Khansa I, Catignani K, et al. Computerized segmentation and measurement of chronic wound images. *Comput Biol Med*, 2015(60):74-85

11 , , , . , 2015, 31(2):86-92

【摘要】目的

方法 22 (22)

11 500ml0.9% 0.5ml1%
500 ml 0.9% 0.1 ml1%

结果 17.238 min

16.179 min (1.71, . . 0.05)

结论 500 ml 0.9%

0.1 ml 1%

【关键词】

DOI 10.14163/j.cnki.11-5547/r.2020.10.048

【摘要】

To discuss the effect of low concentration epinephrine infusion on duration of

ocular anesthesia during cataract extraction combined with intraocular lens implantation. A total

of 22 patients (22 involved eyes) undergoing cataract extraction combined with intraocular lens implantation

were randomly divided into control group and experimental group, with 11 cases in each group. Patients in the

control group received infusion solution consisting of 500 ml of 0.9% sodium chloride solution and 0.5 ml of 1%

adrenaline. Patients in the experimental group received infusion solution consisting of 500 ml of 0.9% sodium

chloride solution and 0.1 ml of 1% adrenaline. The duration of anesthesia during operation was compared between

the two groups. The median maintenance time of anesthesia effect in the control group was 17.238 min,

and the median maintenance time of anesthesia effect in the test group was 16.729 min, and the difference was

not statistically significant (1.71, . . 0.05). In the experimental group, the pupil size at the end of the operation

was equal to that at the beginning of the operation, and it could maintain the state of dilation during the whol-

t